## **COMMUNICABLE DISEASES QUICK REFERENCE**

DISEASE	NOTIFIABLE	HOW CAUSED	INCUBATION PERIOD	INFECTIOUS FOR	PERIOD OF EXCLUSION FROM SCHOOL
CHICKEN POX	NO	Airborne or droplet infection; direct contact with fluid from a vesicle Of an infected person. Once the scabs are dry they are no longer infectious.	10-21 days (usually 13-17 days)	From 2 days before rash until all blisters have crusted.	Exclude until at least 5 days after the eruption first appears. Some remaining scabs do not justify exclusion
CONJUNCTIVITIS	NO	Direct or indirect contact with secretion from infected eyes	1-3 days	While eye discharge is present.	Exclude from school until discharge from eyes has ceased. Contacts: do not exclude.
SCHOOL SORES (IMPETIGO)	NO	Contact with infected person.	1-3 days	Until sores are healed.	Exclude from school until effective treatment (including the proper use of occlusive dressings) has been instituted.
HEAD LICE (PEDICULOSIS)	NO	Contact with infested person or clothing.	1-7 days	Until lice and nits (ova) are destroyed.	Exclude from school until effective treatment has been instituted and nits removed from hair. Contacts: family contacts will probably be infested and should be treated.
INFLUENZA- LIKE ILLNESS	NO	Airborne or droplet infection, or direct contact with contaminated nose or throat secretions.	1-3 days	2 days before onset of symptoms to 5 days after.	Re-admit on recovery.
MEASLES	YES	Spray or droplet or direct contact with contaminated nose or throat secretions.	7-18 days (usually 10-14 days)	4-5 days before to 4 days after the appearance of rash.	Exclude from school. Re-admit on medical certificate of recovery, or at least 4 days from appearance of rash, if well. Contacts: do not exclude.
RINGWORM	NO	Contact with infected persons, articles or animals (especially cats).	4-14 days	As long as lesions contain fungus & spores.	Exclude from school. Re-admit on medical cert. that the child is no longer likely to convey the infection. Contacts: do not exclude.
HEPATITIS A	YES	Contact with case or carrier. Contamination of fingers, foods or drinking water. By faecal material from infected person. Flies may act as vectors.	15-50 days (usually 3 - 5 weeks)	From several weeks before onset of symptoms to 7 days after.	Exclude from school. Re-admit on medical cert. of recovery or on subsidence of symptoms, but not before 7 days after onset of jaundice.